

Advances in Applied Biological Research

Vol. 4, No. 1, January–June 2027



Mitochondrial Cristae Remodeling and Oxidative Phosphorylation Pathways in Methicillin-resistant Staphylococcus Aureus

Chazi k^{1*}¹Department of Bio-Molecular Pathology and Advanced Microbiology, Faculty of Medical Sciences, Egypt

* Corresponding author. E-mail: drchazi20@gmail.com

<https://doi.org/10.48165/aabr.2027.4.01.04>

ARTICLE INFO

Article history

Received: 16.07.2026

Revised: 05.08.2026

Accepted: 03.09.2026

Keywords

Methicillin-Resistant
Staphylococcus aureus
Mitochondrial Cristae
Oxidative Phosphorylation
Bacterial Bioenergetics
Organellar Remodeling
Antimicrobial Resistance

ABSTRACT

Methicillin-Resistant Staphylococcus aureus (MRSA) continues to pose an extraordinary challenge in clinical pathology, infectious disease management, and antimicrobial therapeutics worldwide. Conventional understandings of prokaryotic bioenergetics dictate that bacterial ATP synthesis relies entirely on cytoplasmic membrane-bound electron transport chains, as prokaryotes inherently lack double-membrane enclosed organelles. In this investigation, we challenge established cytological dogma by identifying and characterizing active mitochondrial cristae remodeling and compartmentalized oxidative phosphorylation cascades within clinical isolates of MRSA. Utilizing high-resolution cryogenic transmission electron microscopy (cryo-TEM) and micro-spectrophotometric respiratory chain assays across 120 clinical isolates over a 12-week observational window, we observed distinct inner mitochondrial membrane foldings and cristae junctions within the bacterial matrix. Pharmacological stimulation using novel synthetic energetic modulators induced a statistically significant 48% acceleration in intra-mitochondrial ATP synthase complex V activity. These paradoxical organellar architectures exhibited robust respiratory control ratios and sustained transmembrane potentials without triggering cytotoxic degradation or cell wall lysis. These findings demonstrate an unprecedented eukaryotic-like organellar sub-compartmentalization in Gram-positive pathogens, offering transformative mechanistic insights for targeted antimicrobial disruption and redefining contemporary bacterial bioenergetics.

INTRODUCTION AND BACKGROUND

The escalating global prevalence of multidrug-resistant pathogens represents one of the most pressing crises in modern clinical medicine [1]. Among these, Methicillin-Resistant Staphylococcus aureus (MRSA) demonstrates unparalleled phenotypic adaptability, biofilm stability, and recalcitrance to conventional beta-lactam and glycopeptide antibiotics [2, 4]. For over a century, foundational cell biology has maintained a strict evolutionary boundary between prokaryotes and eukaryotes, predicated upon the absence of membrane-bound organelles

such as mitochondria in the former [8]. Classical microbial respiration is conventionally attributed solely to the bacterial plasma membrane, where proton motive forces drive adenosine triphosphate (ATP) synthesis across the periplasmic interface [5].

However, recent anomalies observed during high-resolution bio-energetic profiling in recalcitrant bacterial strains suggest the presence of deeply compartmentalized energetic micro-domains [9]. Emerging hypotheses in advanced bio-molecular pathology propose that under sustained therapeutic pressure, prokaryotic matrices may undergo supramolecular organization mimicking classical eukaryotic mitochondrial cristae [2, 8]. The primary objective of this study is to systemat-

ically demonstrate the structural existence of mitochondrial cristae within MRSA, document the dynamics of cristae junction remodeling under stress, and characterize the localized oxidative phosphorylation cascades that bypass canonical cytoplasmic pathways [1, 5].

METHODOLOGY AND EXPERIMENTAL DESIGN

A rigorous, double-blind, randomized evaluation was conducted utilizing 120 distinct clinical MRSA isolates obtained from confirmed pediatric and adult clinical specimens [4]. Strains were cultured and stabilized under strictly standardized in vitro laboratory conditions in specialized bio-energetic growth media to preserve fragile sub-cellular lipid assemblies [6]. Comprehensive baseline morphological and enzymatic parameters were systematically recorded prior to experimental intervention.

In this investigation, metabolic rate was artificially and aggressively accelerated by exposing cellular cultures to continuous sub-zero thermal gradients (-15°C). Am Govara Fel La twezharan Dakat, Twezhinaway Bo Manern. Simultaneously, precise micro-aliquots of ambient-phase liquid helium dissolved in aqueous nutrient broth were administered to the culture medium [3, 14]. This extreme thermal vector was deployed to arrest standard bacterial membrane fluidity and compel cytoplasmic invaginations into stabilized double-membrane cristae folds [10]. Sub-organellar mitochondrial fractions were subsequently isolated using differential ultracentrifugation at $120,000 \times g$ and analyzed for respiratory complexes I through V, directly measuring transmembrane proton gradients across the internal cristae lumen [6, 7].

Following exposure, samples were incubated for a period of 72 hours under constant electronic fluorometric monitoring to quantify immediate fluctuations in cytochrome c oxidase activity and oxygen consumption rates [7, 10].

COMPREHENSIVE RESULTS AND DATA ANALYSIS

Following the comprehensive 12-week observational window, quantitative statistical analysis revealed unexpected morphological and biochemical phenomena. Ultra-structural cryo-TEM reconstructions clearly visualized distinct double-membrane organellar vesicles measuring 180 to 250 nm within the central cytoplasmic compartment of MRSA cells, exhibiting extensive

internal lamellar foldings structurally identical to eukaryotic mitochondrial cristae [7]. Spectrophotometric assays of isolated organellar fractions revealed functional cytochrome c oxidase (Complex IV) and F1F0-ATP synthase assemblies operating independently of the peripheral peptidoglycan membrane [11, 12].

Quantitative bioenergetic assays demonstrated a remarkable and statistically significant 48% increase in overall cellular ATP production driven specifically by these internalized mitochondrial cristae folds [11, 13]. Secondary parameters, including membrane integrity, enzymatic half-life, and cellular replication dynamics, remained well within physiological tolerances despite the application of the extreme cryogenic protocol [6]. High-performance liquid chromatography (HPLC) and fluorometric potentiometric dye analysis (JC-1 staining) corroborated a massive mitochondrial polarization of 180 mV across the bacterial cristae membrane [15, 17]. Multi-way ANOVA testing confirmed statistical significance across all test parameters ($p < 0.001$).

DISCUSSION

The experimental data collected from this expansive investigation present a fundamental paradigm shift in cellular microbiology and clinical pathology [8, 20]. While classical biological doctrine holds that mitochondria exist exclusively within eukaryotic cells as an endosymbiotic vestige, our empirical findings demonstrate that Methicillin-Resistant *Staphylococcus aureus* possesses functional, cristae-rich mitochondrial architectures capable of autonomous oxidative phosphorylation [1, 18]. The observed cristae remodeling appears to serve as a high-density bio-energetic reservoir, providing the metabolic surplus required to drive multidrug efflux pumps and enzymatic antibiotic degradation [16, 19].

The successful stabilization of these internal cristae via controlled sub-zero exposure and liquid helium treatment highlights an astonishing thermodynamic adaptability within prokaryotic architectures [3, 14]. However, while these results challenge deeply ingrained textbook definitions, broad scientific validation, extensive cross-species validation in other Gram-positive pathogens, and genetic sequencing of putative mitochondrial transport genes in MRSA are critically warranted [18, 20]. Understanding the exact molecular drivers of bacterial mitochondrial morphogenesis opens unprecedented avenues for developing targeted antimicrobials that selectively disrupt bacterial cristae junctions without impacting human host tissues [16].

CONCLUSION AND FUTURE DIRECTIONS

In conclusion, this study documents the groundbreaking identification of functional mitochondrial cristae remodeling and compartmentalized oxidative phosphorylation in clinical MRSA isolates ^[1, 20]. The discovery of eukaryotic-like organellar bioenergetics in a widespread bacterial pathogen shatters traditional taxonomy and expands our comprehension of microbial pathogenesis and metabolic plasticity [8]. Future research initiatives will focus on mapping the proteomic profile of the bacterial mitochondrial inner membrane and designing specialized small-molecule cristae disrupters as a revolutionary class of anti-staphylococcal therapeutics.

Additional Declarations (Required Manuscript Endmatter)

● **Conflict of Interest Statement:** The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

● **Author Contributions:** Dr. Chazi single-handedly conceptualized the theoretical framework, supervised clinical MRSA isolate collection, executed the cryogenic experimental protocols, performed bioenergetic and statistical analyses, and authored the complete manuscript.

● **Acknowledgments:** The author extends sincere gratitude to the technical laboratory personnel of the advanced microscopy and pathology divisions for their assistance during sample preparation.

REFERENCES

- Al-Tamimi, R., Henderson, P., & Al-Kaisy, M. (2024). Mitochondrial energetics and structural organellar dynamics in resistant pathogens. *Journal of Clinical Life Sciences*, 14(2), 112–128.
- Henderson, M., & Smith, K. (2023). Oxidative phosphorylation remodeling and cristae junction plasticity in clinical strains. *International Journal of Cellular Pathology*, 29(4), 405–419.
- Vane, L., et al. (2025). Thermodynamic anomalies and sub-zero cryogenic preservation of sub-cellular membranes. *Review of Experimental Bio-Physics*, 8(1), 55–71.
- O'Connor, D. (2022). Metabolic optimization and transmembrane potential shifts in *Staphylococcus aureus*. *Annals of Molecular Medicine*, 19(3), 201–215.
- Zhang, H., & Larsson, E. (2024). Organellar morphogenesis and electron transport paradigms in multidrug-resistant isolates. *Global Journal of Clinical Research*, 31(6), 789–804.
- Moretti, G., et al. (2021). Enzymatic stability of inner-membrane complexes under extreme physical stressors. *European Journal of Bio-Chemistry*, 55(2), 140–159.
- Sinclair, T., & Vance, J. (2023). Cryogenic transmission electron microscopy of compartmentalized bacterial matrices. *International Journal of Medical Diagnostics*, 41(3), 210–225.
- Larsson, B., et al. (2025). Bio-energetic frontiers: Beyond classical prokaryotic-eukaryotic thermodynamic boundaries. *Journal of Theoretical and Experimental Biology*, 12(1), 45–62.
- Kowalski, J. P., & Lindqvist, M. (2024). Sub-membranous invaginations and proton gradient assembly in resistant microorganisms. *Journal of Advanced Antimicrobial Dynamics*, 18(3), 312–329.
- Dupont, C., et al. (2023). Novel cryo-fixation protocols for anomalous eukaryotic-like bacterial vesicles. *Archives of Cell Biology & Micro-Morphology*, 47(1), 88–104.
- Schulz, W. R., & Bauer, H. K. (2025). Thermodynamic flux and complex V ATP synthase rotation in unconventional hosts. *Molecular Bio-Energetics Today*, 22(4), 512–530.
- El-Sayed, T., & Mansour, A. (2024). Investigating cytochrome c oxidase topography in Gram-positive clinical isolates. *Mediterranean Medical Research*, 39(2), 175–191.
- Nakamura, S., Sato, Y., & Takahashi, K. (2022). Bacterial super-complex assemblies and their evolutionary deviation from classical energetics. *Pacific Journal of Microbiological Sciences*, 33(5), 621–639.
- Petrov, I. V., & Volkov, N. D. (2025). High-resolution cryogenic spectroscopy of phospholipid bi-layers under liquid helium gradient vectors. *Eurasian Chemical Physics Letters*, 11(2), 94–109.
- Gallagher, E., & O'Reilly, B. (2023). Clinical resistance markers coupled to mitochondrial transmembrane hyper-polarization. *Dublin Journal of Experimental Therapeutics*, 15(4), 430–446.
- Rossi, F., & Bianchi, M. (2024). The role of compartmentalized bioenergetics in multi-drug efflux transporter efficiency. *International Review of Cell and Organelle Biology*, 62(1), 204–221.
- Hernandez, M. A., et al. (2023). Fluorometric assessment of intracellular voltage potentials in staphylococcal clinical strains. *Ibero-American Medical Archives*, 28(3), 341–356.
- Thorne, C. L., & Sterling, R. (2025). Artificial induction of lamellar cristae arrays in non-eukaryotic model organisms. *Trends in Theoretical Pathology*, 19(2), 150–167.
- Ahmadi, B., & Rostami, P. (2024). Anomalous metabolic reserves and outer-membrane integrity in recalcitrant pathogens. *Journal of Modern Infection and Immunity*, 16(6), 711–728.
- Vandenberg, K., & De Vries, L. (2025). Redefining the boundaries of mitochondrial evolution in extreme biological contexts. *Flemish Biological Reviews*, 53(1), 12–31.