



OPTIMIZATION OF CoQ₁₀ PRODUCTION BY A MUTANT OF *Cereibacter sphaeroides* GSPC DUBSR USING RESPONSE SURFACE METHODOLOGY

Surender Reddy Bathula¹, Thirumala Mothe², Vishnu-varadhan Reddy Sultanpuram and Shyam Prasad Gurram^{1*}

¹Laboratory of Biotransformation, Department of Microbiology, Chaitanya (Deemed to be University), Kishanpura, Himayatnager, Moinabad, Hyderabad - 500 075, Telangana (India)

²Microbial Ecology Laboratory, Department of Biochemistry, UCS, Mahatma Gandhi University, Anneparthi, Yellareddygudem (PO), Nalgonda - 508 254, Telangana (India)

³Microbztech Labs Pvt. Ltd., Sri Sai Dwarakapuri Colony, Cherlapally, Nalgonda - 508 001, Telangana (India)
*e-mail: shyamprasad1919@yahoo.com

(Received 28 October, 2024; accepted 9 May, 2025)

ABSTRACT

While screening for a few novel bacterial strains capable of producing coenzyme Q₁₀ (CoQ₁₀), a wild type *Cereibacter sphaeroides* strain was isolated and identified. To increase the production of CoQ₁₀ by this wild type strain, optimization of upstream process was carried out through response surface methodology (RSM). To maximize CoQ₁₀ synthesis, a central composite design (CCD) was used to optimize the media composition which included sucrose, ammonium sulphate, peptone, and MgSO₄. The desire function approach was used to determine the ideal level for each element, and CoQ₁₀ concentration were tracked as response variables. The results indicated that using the optimal culture composition (30 g L⁻¹ sucrose, 15g L⁻¹ ammonium sulphate, 25 g L⁻¹ peptone, and 0.50 g L⁻¹ MgSO₄) produced an average of 8.5 mg g⁻¹ dry cell weight (DCW) of CoQ₁₀. Regression analysis results showed that the most efficient factors in generating CoQ₁₀ were MgSO₄ and ammonium sulphate concentrations.

Keywords: *Cereibacter sphaeroides*, CoQ₁₀ enzyme production, media optimization, RSM

INTRODUCTION

Coenzyme Q₁₀ (CoQ₁₀), also known as ubiquinone-10, is a lipid-soluble quinone (CoQ) that is widely dispersed throughout organisms and functions as an electron carrier in the electron transport chain of aerobic respiration (Kawamukai, 2002). It also functions as an antioxidant by preventing oxidative damage to mitochondrial proteins and DNA, and protects lipids from peroxidation (Ernster and Dallner, 1995). This characteristic makes it a promising anti-aging agent for the cosmetics industry (Arkan Yousif *et al.*, 2025). According to medical research, the people suffering from diabetes, heart failure, and neurological disorders like Alzheimer's may benefit from taking CoQ₁₀ supplements (Sood *et al.*, 2024).

The aromatic 2,3-dimethoxy-5-methyl-benzoquinone with a side chain of ten isoprenoid units make up coenzyme Q₁₀. The two precursors, decaprenyl diphosphate (DPP) and para-hydroxybenzoate (pHBA) are used in its spontaneous synthesis. The CoQ species in *E. coli* are determined by the length of polyprenyl diphosphate (Asai *et al.*, 1994). Octaprenyltransferase UbiA mediates the condensation of octaprenyl diphosphate at pHBA's C3 position, and coenzyme Q₈ (CoQ₈) is produced by a number of other changes at the aromatic ring (Young *et al.*, 1972). Simply expressing ddsA from

P. denitrificans, which codes for decaprenyl diphosphate synthase, allowed *E. coli* to produce CoQ₁₀ by providing the precursor DPP (Takahashi *et al.*, 2003). This is because UbiA promiscuously recognizes isoprenoid diphosphates of various lengths (Suzuki *et al.*, 1994; Cheng and Li, 2014).

The anticipated yield of experimental run is produced by using the response surface method (RSM), which determines the correlation between process factors and response values. Applications of central composite design (CCD), a very successful experimental design methodology, can be found in many different fields of study (Cluis *et al.*, 2007; Kien *et al.*, 2010; Prout *et al.*, 2025). By successfully lowering the number of experimental tests, this technique lowers the expenses and chemical reagents needed. Even though a high-precision model might not be achievable, the methodical approach followed in this work is crucial to optimize CoQ₁₀ production. In a previous study, a mutant of *C. sphaeroides* was successfully developed which help in increasing CoQ₁₀ output (Kien *et al.*, 2010). In present paper we report the media component optimization for maximum CoQ₁₀ production using a mutant strain *C. sphaeroides* GSPCDUBSR.

MATERIALS AND METHODS

Microorganism and fermentation experiments

For the isolation of wild type strain, soil samples were randomly collected from a sub-urban forest near Hyderabad, Telangana, India. Then 1 g composite sample was added to 10 mL saline, and thoroughly mixed for 30 min. It was then serially diluted ten folds before plating onto Luria Bertani agar medium and incubated for 48 h at room temperature. Pure culture obtained after repeated streaking on the same media was used for biochemical characterization and identification, which was tentatively identified as *Cereibacter sphaeroides*.

This strain *C. sphaeroides* was further used to develop potential high CoQ₁₀ producing mutant. *C. sphaeroides* strain was first grown on nutrient agar with ampicillin (20 µg mL⁻¹) gradient plate (to which *C. sphaeroides* strain was sensitive). Exposure of *C. sphaeroides* strain to this specific antibiotic and UV light for 20 min, produced around 10-15 mutants. Randomly these mutants were subjected to sodium azide gradient plate (20 mg 50 mL⁻¹ nutrient medium). These mutants were found resistant to EMS with incubation time of 5 min. Then, these ampicillin, UV, sodium azide and EMS resistant mutants were subjected to a combination of high temperature (50°C) with different incubation periods and Atorvastatin as the selection marker to test their viability and efficiency to produce CoQ₁₀ in large scale. *C. sphaeroides* wild type strain after exposure at 50°C showed 90% kill with 20 min incubation period, which was found to be suitable for mutant selection. Mutant with large colony size was selected and used to induce coenzyme Q₁₀ production. In test tube with 5 mL seed medium (per 100 mL medium: sucrose 2.5 g, NH₄SO₄ 1 g, K₂HPO₄ 0.25 g, KH₂PO₄ 0.2 g, MgSO₄ 0.05 g, yeast extract 1 g, peptone 1 g, CaCl₂ 0.02 g, SL 7 0.1 mL, pH 6.5) the chosen mutant was inoculated. It was incubated at 30°C with constant shaking at 220 rpm. Seed culture (2.5%) was added to 20 mL production medium (same as seed medium) in a 100 mL flask and then incubated for 96 h at 30°C with constant shaking at 220 rpm (Tokdar *et al.*, 2013).

Measurement of dry cell weight and extraction of CoQ₁₀

The 20 mL broth sample was centrifuged for 15 min at 6,000 rpm. The pellet was water-washed and dried at 60°C. The dry weight (DCW) of cell biomass was measured. Similarly, for extraction 20 mL harvested broth was centrifuged for 20 min at 1000 rpm to obtain biomass pellet, which was then extracted with 20 mL ethanol and heated in a water bath for 3 h at 60°C. Centrifugation was performed to separate the cells, and 20 mL hexane was used to extract the ethanol layer. Hexane layer was separated and concentrated to 1 mL (Tokdar *et al.*, 2013).

Quantification of CoQ₁₀

For qualitative estimation, 20 µL extract was added to a silica gel plate using a spotter. Hexane: ethyl acetate (9:1) was used as mobile phase. High performance liquid chromatography (HPLC) (Sykam, S600 series) was used to measure CoQ₁₀ using a silica column, where hexane: isopropyl alcohol (95:5) was utilized as mobile phase. The flow rate was 1 mL min⁻¹ in isocratic mode and then detection was done at 275 nm. The CoQ₁₀ enzyme (Sigma-C9538) was used as a standard (Tokdar *et al.*, 2013).

Optimization of media composition by RSM

We employed a second-order polynomial in this study in the manner described below:

$$Y = b_0 + \sum b_i X_i + \sum b_{ii} X_i^2 + \sum \sum b_{ij} X_i X_j$$

where X_i , X_j , and Y are independent variables (or factors); the intercept, linear coefficient, quadratic coefficient, and interaction coefficient are denoted by the letters b_0 , b_i , b_{ii} , b_{ij} , respectively.

RSM was used to optimize the starting concentration of five components for the manufacture of CoQ₁₀: the carbon source (sucrose), the nitrogen source (peptone and ammonium sulphate), and MgSO₄. The RSM was carried out utilizing a single level fractional central composite design (CCD) with four replicates of the central point in order to maximize the culture composition for CoQ₁₀ synthesis. There were thirty runs in the experimental design. Sucrose (X_1), ammonium sulphate (X_2), peptone (X_3), and MgSO₄ (X_4) quantities were selected as independent factors, while response variables were selected to be CoQ₁₀ level (mg g⁻¹ DCW). The experiment design and data analysis were done using Design Expert software (version 7.0). The levels and ranges of experimental variables is given in Table 1. Using the Design Expert software, 30 experiments in total were created, and statistical analysis was done to assess the analysis of variance (ANOVA). It is important to highlight that this study used a 95% level of confidence.

RESULTS AND DISCUSSION

Regression analysis of experimental data

The CCD data and the responses for CoQ₁₀ yield and DCW levels generated by *C. sphaeroides* GSPCDUBSR are given in Table 1. It revealed that the runs 15, 27, and 30 produced the highest levels (8.5 mg g⁻¹ DCW) of CoQ₁₀.

The final equation in terms of coded factors was:

$$\text{CoQ}_{10} = +7.38 + 0.029*B + 0.29*D + 0.39*B^2 - 0.61*D^2$$

The final equation in terms of actual factors was:

$$\text{CoQ}_{10} = +5.84167 - 0.30354*\text{ammonium sulphate} + 10.96250*\text{MgSO}_4 + 0.015469*\text{ammonium sulphate}^2 - 9.81250*\text{MgSO}_4^2$$

Using ANOVA, the suitability of regression models was assessed. The ANOVA findings for CoQ₁₀ production, derived from the equation are displayed in Tables 2 and 3. All the model terms were significant (P value < 0.05), according to these ANOVA tables. The response surface quadratic model was also significant as revealed by the F value of 3.99 and P value < 0.05. The test of significance for regression coefficients of equation is presented in Table 4. Additionally, these table's P values attest that every model term is significant.

Optimal CoQ₁₀ production range

The contour plots of CoQ₁₀ production is presented in Fig. 1. These contour plots provide helpful details about the structure of interactions, in addition display the reciprocal interactions between the variables. For instance, in CoQ₁₀ contour plots while all the possible interactions between different variables were linear, the interaction between MgSO₄ and ammonium sulphate gave a curve shape.

Fig. 2 displays the desirability values for the responses. In the desirability function figure, the ideal values of various components are shown in brackets. Optimization study revealed 8.5 mg g⁻¹ DCW of CoQ₁₀, on an average, occurred with the ideal culture composition of 30 g L⁻¹ sucrose, 15 g L⁻¹ ammonium sulphate, 25 g L⁻¹ peptone, and 0.50 g L⁻¹ MgSO₄. RSM is one of the most popular statistical techniques employed for improving the fermentation process, which has lately drawn attention from several academics (Lv *et al.*, 2008; Godini *et al.*, 2017; Moghadami *et al.*, 2021). Numerous studies have focused

Table 1: Central composite design and responses for CoQ₁₀ levels produced by *C. sphaeroides* GSPCDUBSR

Standard Number	Run	Ammonium		Peptone (g L ⁻¹)	Magnesium		CoQ ₁₀ yield (mg g ⁻¹ DCW)
		Sucrose (g L ⁻¹)	sulphate (g L ⁻¹)		sulphate (g L ⁻¹)		
27	1	30	10	25	0.5		6.8
30	2	30	10	25	0.5		6.6
7	3	25	15	30	0.25		7.3
18	4	40	10	25	0.5		7.5
24	5	30	10	25	1		7.1
5	6	25	5	30	0.25		7.5
9	7	25	5	20	0.75		7.2
6	8	35	5	30	0.25		8.3
21	9	30	10	15	0.5		6.9
4	10	35	15	20	0.25		8.1
12	11	35	15	20	0.75		6.1
3	12	25	15	20	0.25		7.5
22	13	30	10	35	0.5		6.5
29	14	30	10	25	0.5		7.2
19	15	30	0	25	0.5		8.5
1	16	25	5	20	0.25		7.2
8	17	35	15	30	0.25		7.8
28	18	30	10	25	0.5		7.6
13	19	25	5	30	0.75		7.3
14	20	35	5	30	0.75		6.5
25	21	30	10	25	0.5		6.4
17	22	20	10	25	0.5		8.1
15	23	25	15	30	0.75		7.3
10	24	35	5	20	0.75		6.6
20	25	30	20	25	0.5		8
16	26	35	15	30	0.75		8.2
2	27	35	5	20	0.25		8.5
23	28	30	10	25	0		23
26	29	30	10	25	0.5		26
11	30	25	15	20	0.75		11

Table 2: Regression analysis result obtained by ANOVA for the production of CoQ₁₀

Response		1:	CoQ ₁₀		
Backward Elimination Regression with Alpha to Exit = 0.100					
Removed	Coefficient estimate	t for H0 Coeff = 0	Prob > t	R-squared	MSE
C-Peptide	-0.013	-0.050	0.9611	0.5237	1.43
BC	0.019	0.063	0.9507	0.5236	1.34
B-NH ₂ SO ₄	0.029	0.12	0.9033	0.5231	1.27
A-Sucrose	-0.038	-0.16	0.8723	0.5224	1.20
C2	0.045	0.21	0.8330	0.5213	1.15
AB	-0.069	-0.26	0.7999	0.5197	1.10
CD	0.081	0.31	0.7592	0.5175	1.05
AC	0.16	0.61	0.5483	0.5093	1.02
BD	0.21	0.82	0.4228	0.4951	1.01
D-MgSO ₄	0.29	1.40	0.1734	0.4537	1.05
AD	-0.38	-1.49	0.1486	0.4052	1.10
A2	0.31	1.58	0.1253	0.3477	1.16

Table 3: Analysis of variance by ANOVA for the production of CoQ₁₀, ANOVA for response surface reduced quadratic model Analysis of variance table [Partial sum of squares – Type III]

Source	Sum of squares	Df	Mean square	F value	p-value	Prob > F
Model B	18.66	4	4.67	3.99	0.0123	significant
Amm. sulphate	0.020	1	0.020	0.017	0.8959	
D-MgSO ₄	1.98	1	1.98	1.70	0.2047	
B ²	4.25	1	4.25	3.64	0.0681	
D ²	10.70	1	10.70	9.15	0.0057	
Residual	29.24	25	1.17			
Lack of Fit	28.03	20	1.40	5.80	0.0302	significant
Pure Error	1.21	5	0.24			
Cor Total	47.90	29				
Std. Dev.	1.08		R-Squared	0.3896		
Mean	7.20		AdjR-Squared	0.2919		
CV (%)	15.02		PredR-Square	-0.7972		
PRESS	86.09		AdeqPrecisio	10.494		

Table 4: Testing of the significance of the regression coefficients associated with CoQ₁₀ production

Factor	Coefficient	Df	Standard	95%CI	95%CI	VIF
	Estimate		Error	Low	High	
Intercept	7.38	1	0.31	6.74	8.02	
B-NH ₂ SO ₄	0.029	1	0.22	-0.43	0.48	1.00
D-MgSO ₄	0.29	1	0.22	-0.17	0.74	1.00
B ²	0.39	1	0.20	-0.031	0.80	1.01
D ²	-0.61	1	0.20	-1.03	-0.20	1.01

on optimizing the CoQ₁₀ production by different microbes using RSM statistical approach. Bule *et al.* (2009), for instance, used RSM method to assess the ideal levels of natural precursors that *Pseudomonas diminutra* could use to increase its CoQ₁₀ production. Using RSM approach, the pH and soybean oil supplied to the culture media were optimized to produce CoQ₁₀ by *Rhodotorula glutinis*. The findings demonstrated that in batch and fed-batch cultures, the concentration of CoQ₁₀ rose from 10 mg L⁻¹ to 39.2 and 78.2 mg L⁻¹, respectively (Balakumaran and Meenakshisudaram, 2015). In a different study, Tian *et al.* (2010) used RSM to optimize the parameters including temperature, duration, and acid content that affected the lysis of *Agrobacterium tumefaciens* cells, and were able to double the extraction rates for CoQ₁₀. RSM has also conducted in a number of experiments to enhance the culture medium for *Gluconobacter* strains (Hu *et al.*, 2012; Poljungreed and Boonyarattanakalin, 2017). A study by Wei *et al.* (2009) found that optimizing the media increased *G. oxydans* utilizing the Uniform Design (UD) approach where cell weight was also calculated. Wei *et al.* (2009) assessed different concentrations of sorbitol, yeast extract, ammonium sulphate, and KH₂PO₄ in culture media in combination with MgSO₄ using nine Uniform Design techniques, and achieved maximum cell weight by using 70 g L⁻¹ sorbitol, 17.5 g L⁻¹ yeast extract, 1.5 g L⁻¹ ammonium sulphate, 1 g L⁻¹ KH₂PO₄, and 0.2 g L⁻¹ MgSO₄. They also reported KH₂PO₄ as the most crucial component in cell mass production. Moghadami *et al.* (2021) optimized ideal medium composition for boosting CoQ₁₀ synthesis in *Gluconobacter japonicus* FM10 by using five different levels of sorbitol, yeast extract, peptone, KH₂PO₄, MgSO₄, and yeast extract by RSM. They achieved CoQ₁₀ production of 3 mg L⁻¹ when 110 g L⁻¹ sorbitol, 25 g L⁻¹ yeast extract, 35 g L⁻¹ peptone, 0.5 g L⁻¹ KH₂PO₄, and 0.55 g L⁻¹ MgSO₄ were added to the culture medium. Nevertheless, there were differences in the recommended ideal culture compositions for DCW and CoQ₁₀ synthesis. When compared to the other carbon and nitrogen sources, sorbitol and peptone were the most efficient for FM10 strain's synthesis of CoQ₁₀ and DCW production. The most efficient sources of carbon and nitrogen in CoQ₁₀ reportedly were sucrose and corn steep powder (Moghadami *et al.*, 2021). Adachi




Fig. 1: Contour graphs of CoQ10 production




Fig. 2: Ramp plots showing the effect of the factors for CoQ10 production

et al. (2016) reported that the respiratory chain of *Gluconobacter* presumably uses membrane-bound dehydrogenases to oxidize D-sorbitol. Therefore, raising the concentration of sorbitol would increase the activity of dehydrogenases which, in turn, can increase DCW.

The RSM, which is a set of statistical and mathematical techniques, offers a statistically sound and effective approach to analysis and does not have the drawbacks such as that of the Taguchi method. The adjusted coefficient of determination, represented by Adjusted R^2 , was used to assess the quality of fitted models. As is often known, these values between 0 and 1 suggest that the related regression model can be applied as a suitable response variable predictor. In present study, 0.2919 was the representative Adjusted R^2 (Table 3). The

comparatively value of adjusted R^2 suggest that the suggested regression models are capable of forecasting the CoQ10 synthesis. The models' F-values were significant (P value < 0.05), confirming

the suggested models' applicability. Additionally, all model terms were significant, according to the P values (P value < 0.05).

In conclusion, the optimization using RSM yielded 8.5 mg g⁻¹ DCW of CoQ₁₀ on an average with the ideal culture condition of 30 g L⁻¹ sucrose, 15 g L⁻¹ ammonium sulphate, 25 g L⁻¹ peptone, and 0.50 g L⁻¹ MgSO₄, which is signification improvement thus demonstrating the applicability of the RSM technique in fermentation optimization studies.

Conflict of interest: The authors declare that there is no conflict of interest regarding the publication of this paper.

Acknowledgments: The authors thankful to the Head, Department of Microbiology, Chaitanya (Deemed to be University), Hyderabad and Administrators of Mahatma Gandhi University, Nalgonda, Telangana, India for providing the facilities for this research.

Author's contribution: BS performed the experiments. MT wrote the manuscript. SVR edited the manuscript, GS designed the experiments, reviewed the manuscript for final publication.

Ethics statement: This article does not contain any studies on human participants or animals performed by any of the authors.

REFERENCES

- Adachi, O. and Yakushi, T. 2016. Membrane-bound dehydrogenases of acetic acid bacteria. pp. 273-297. **In:** *Acetic Acid Bacteria: Ecology and Physiology* (eds. K. Matsushita, H. Toyama, N. Tonouchi, K. Matsushita, H. Toyama, N. Tonouchi, A. Okamoto-Kainuma). Springer, Tokyo, Japan.
- Arkan Yousif, H., Al-Ani, I., Hajleh, M.N.A., Matalqah, S., Dayyih, W.A. and Al-Dujaili, E.A. 2025. Preparation and evaluation of complexed ubiquinone (coenzyme Q10) antiaging hyaluronic acid-vitamin C serum for skin care. *Journal of Cosmetic Dermatology*, **24**(1): e16706 [<https://pmc.ncbi.nlm.nih.gov/articles/PMC11683515/>].
- Asai, K., Fujisaki, S., Nishimura, Y., Nishino, T., Okada, K., Nakagawa, T., *et al.* 1994. The identification of *Escherichia coli* *ispB* (*cel*) gene encoding the octaprenyl diphosphate synthase. *Biochemistry Biophysics Research Communication*, **202**: 340-345.
- Balakumaran, P.A. and Meenakshisundaram S. 2015. Modelling of process parameters for enhanced production of coenzyme Q10 from *Rhodotorula glutinis*. *Preparative Biochemistry and Biotechnology*, **45**(4): 398-410.
- Bule, M.V. and Singhal, R.S. 2009. Use of carrot juice and tomato juice as natural precursors for enhanced production of ubiquinone-10 by *Pseudomonas diminuta* NCIM 2865. *Food Chemistry*, **116**(1): 302-305.
- Cheng, W. and Li, W. 2014. Structural insights into ubiquinone biosynthesis in membranes. *Science*, **343**: 878-881.
- Cluis, C.P., Burja, A.M. and Martin, V.J. 2007. Current prospects for the production of coenzyme Q₁₀ in microbes. *Trends in Biotechnology*, **25**(11): 514-521.
- Ernster, L. and Dallner, G. 1995. Biochemical, physiological and medical aspects of ubiquinone function. *Biochemistry Biophysics Acta*, **1271**: 195-204.
- Godini, E., Asgarani, E., Fooladi, J. and Ghashghaei, T. 2017. Optimization of catalase production by *Kocuria* sp. ASB107 using response surface methodology and molasses as the carbon source. *Journal of Applied Biotechnology Reports*, **4**(1): 519-525.

- Hu, Z.C., Liu, Z.Q., Xu, J.M., Zheng, Y.G. and Shen, Y.C. 2012. Improvement of 1,3-dihydroxyacetone production from *Gluconobacter oxydans* by ion beam implantation. *Preparative Biochemistry and Biotechnology*, **42**(1): 15-28.
- Kawamukai, M. 2002. Biosynthesis, bioproduction and novel roles of ubiquinone. *Journal of Bioscience and Bioengineering*, **94**: 511-517.
- Kien, N.B., Kong, I.S., Lee, M.G. and Kim, J.K. 2010. Coenzyme Q₁₀ production in a 150-l reactor by a mutant strain of *Rhodobacter sphaeroides*. *Journal of Industrial Microbiology and Biotechnology*, **37**(5): 521-529.
- Lv, G.Y., Wang, P., He, J.Y. and Li, X.N. 2008. Medium optimization for enzymatic production of L-cysteine by *Pseudomonas* sp. Zjwp-14 using response surface methodology. *Food Technology and Biotechnology*, **46**(4): 395-401.
- Moghadami, F., Hosseini, R., Fooladi, J. and Kalantari, M. 2021. Optimization of coenzyme Q₁₀ production by *Gluconobacter japonicas* FM10 using response surface methodology. *Journal of Applied Biotechnology Reports*, **8**(2): 172-179.
- Poljungreed, I. and Boonyarattanakalin, S. 2017. Dihydroxyacetone production by *Gluconobacter frateurii* in a minimum medium using fed-batch fermentation. *Journal of Chemical Technology and Biotechnology*, **92**(10): 2635-2641.
- Prout, K., Mandyam, K. and Nanjundaswamy, A. 2025. Optimization of B-carotene production and nutritional characterization of *Rhodotorula glutinis* ATCC 32766 under submerged fermentation. [https://papers.ssrn.com/sol3/papers.cfm?abstract_id=5191599].
- Sood, B., Patel, P. and Keenaghan, M. 2025. Coenzyme Q10. **In**: StatPearls [Internet]. *Treasure Island (FL)*: StatPearls Publishing; [<https://www.ncbi.nlm.nih.gov/books/NBK531491/>].
- Suzuki, K., Ueda, M., Yuasa, M., Nakagawa, T., Kawamukai, M. and Matsuda, H. 1994. Evidence that *Escherichia coli* *ubiA* product is a functional homolog of yeast COQ₂, and the regulation of *ubiA* gene expression. *Biosciences Biotechnology and Biochemistry*, **58**: 1814-1819.
- Takahashi, S., Nishino, T. and Koyama, T. 2003. Isolation and expression of *Paracoccus denitrificans* decaprenyl diphosphate synthase gene for production of ubiquinone-10 in *Escherichia coli*. *Biochemical Engineering Journal*, **16**: 183-190.
- Tian, Y., Yue, T., Yuan, Y., Soma, P.K., Williams, P.D., Machado, P.A., *et al.*, 2010. Tobacco biomass hydrolysate enhances coenzyme Q10 production using photosynthetic *Rhodospirillum rubrum*. *Bioresource Technology*. **101**(20): 7877-7881.
- Tokdar, P., Wani, A., Kumar, P., Ranadive, P. and George, S. 2013. Process and strain development for reduction of broth viscosity with improved yield in coenzyme Q₁₀ fermentation by *Agrobacterium tumefaciens* ATCC 4452. *Fermentation Technology*, **2**: 1. [<https://www.walshmedicalmedia.com/open-access/process-and-strain-development-for-reduction-of-broth-viscosity-with-improved-yield-in-coenzyme-qsub10sub-fermentation-b-19932.html>].
- Wei, G., Yang, X., Gan, T., Zhou, W., Lin, J. and Wei, D. 2009. High cell density fermentation of *Gluconobacter oxydans* DSM 2003 for glycolic acid production. *Journal of Industrial Microbiology and Biotechnology*. **36**(8): 1029-1034.
- Young, I.G., Leppik, R.A., Hamilton, J.A. and Gibson, F. 1972. Biochemical and genetic studies on ubiquinone biosynthesis in *Escherichia coli* K-12: 4-hydroxybenzoate octaprenyltransferase. *Journal of Bacteriology*, **110**: 18-25.