Review article

Online ISSN : 3107-9938

Journal of Veterinary and Life Sciences

Journal homepage: www.acspublisher.com

Year 2026, Volume-2, Issue-1 (January–June)

Quorum Sensing and Quenching: Emerging Frontiers in Antimicrobial Therapy Against Drug-resistant Pathogens

Samiksha Sharma[1], Pawan Kumar Verma[2*], Manish Kumar Verma[3] and S R Upadhyay[4]

[1]*MVSc Scholar, Division of Veterinary Pharmacology and Toxicology, FVSc & AH, R S Pura, Jammu, 181102, J&K, India*
[2]*Professor, Division of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Science and Animal Husbandry, R S Pura, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, 181102, India*
[3]*Assistant Professor, Veterinary Pharmacology and Toxicology, Sanskaram College of Veterinary and Animal Science, Jhajjar, Haryana-124108, India*
[4]*Professor, Division of Veterinary Medicine, FVSc & AH, R S Pura, Jammu, 181102, J&K, India*

Keywords:

Antimicrobial resistance
Autoinducers
Biofilm formation
Gene regulation
Quorum sensing

ABSTRACT

Quorum sensing (QS) is a sophisticated cell-to-cell communication mechanism employed by bacteria to regulate gene expression in response to fluctuations in population density. This regulatory system relies on the synthesis, secretion, accumulation, and detection of signalling molecules known as autoinducers. In Gram-negative bacteria, acyl homoserine lactones (AHLs) serve as the primary autoinducers, whereas Gram-positive bacteria predominantly utilize processed oligopeptides as signalling molecules. As bacterial populations expand, these molecules accumulate in the surrounding environment and, upon reaching a critical threshold concentration, trigger coordinated gene expression across the microbial community. This collective response governs a variety of crucial physiological and pathogenic behaviours, including biofilm formation, virulence factor production, bioluminescence, motility, and antimicrobial resistance. In recent years, quorum sensing inhibition (QSI) has emerged as an important and rapidly advancing area of research. Unlike conventional antimicrobial agents that directly target bacterial viability, QSIs disrupt bacterial communication networks, thereby attenuating pathogenicity without exerting strong selective pressure for the development of resistance. By interfering with key stages of quorum sensing pathways, QSIs effectively reduce virulence expression, inhibit biofilm development, and impair bacterial adaptation mechanisms. Consequently, quorum sensing and its inhibition represent promising avenues for the development of innovative antimicrobial therapies and next-generation therapeutic interventions aimed at combating the growing global threat of antimicrobial resistance.

Introduction

The discovery of antimicrobial agents has revolutionized modern medicine and has significantly contributed to the control and treatment of infectious diseases. However, the widespread and often indiscriminate use of these drugs has accelerated the emergence and dissemination of antimicrobial-resistant microorganisms, creating a major One Health challenge with profound implications for human, animal, and environmental health. With bacterial infections accounting for nearly 16 million human deaths annually, there is an urgent need to identify and develop innovative strategies capable of combating drug-resistant pathogens and the diseases associated with them. Extensive investigations into antimicrobial resistance (AMR) have revealed that one of the major contributors to persistent infections and treatment failure is the formation of bacterial biofilms (Musk and Hergenrother, 2006). In addition, numerous bacterial be-

How to cite: Sharma S., Verma P.K., Verma M.K., Upadhyay S.R. (2026). Quorum Sensing and Quenching: Emerging Frontiers in Antimicrobial Therapy Against Drug-resistant Pathogens. *Journal of Veterinary and Life Sciences*, ***2***(1), 21-30. https://doi.org/10.48165/jvls.2026.2.01.04

[*] **Corresponding author E-mail addresses:** drpawankv@yahoo.co.in (Pawan Kumar Verma)
DOI: https://doi.org/10.48165/jvls.2026.2.01.04
Received 01-06-2026, Revision 19-06-2026, Accepted 22-06-2026

Published by ACS Publisher. This article is published under the CC BY-NC license (https://creativecommons.org/licenses/by-nc/4.0/).

haviours associated with pathogenicity and survival are regulated by a phenomenon known as quorum sensing (QS) (Lewis, 2007). QS is a highly coordinated mechanism of microbial communication that enables microorganisms to monitor and respond to changes in their population density. This phenomenon is widely distributed in both Gram-negative and Gram-positive bacteria and has also been documented in diverse microbial groups, including archaebacteria, obligate anaerobes, fungi, and certain eukaryotic hosts. In addition, QS plays an important role in plant–microbe interactions, particularly during root nodulation by nitrogen-fixing bacteria. Interestingly, analogous communication systems resembling QS have also been observed in several social insect species, where they facilitate collective decision-making processes.

QS involves the production, release, and detection of chemical signalling molecules known as autoinducers (AIs). When the concentration of these molecules reaches a specific threshold level, a synchronized gene expression response is initiated throughout the bacterial population (Rutherford and Bassler, 2012). Through this mechanism, bacteria regulate a broad spectrum of physiological activities, including the production of virulence factors such as lectin, exotoxin A, pyocyanin, and elastase in *Pseudomonas aeruginosa* during growth and infection (Antunes et al., 2010), motility (Bramhachari et al., 2018), sporogenesis (Li et al., 2011), bioluminescence (Popham and Stevens, 2006), and biofilm formation (Preda and Sandulescu, 2019). Likewise, the synthesis and secretion of enzymes and virulence determinants such as hemolysins, protein A, enterotoxins, lipases, and fibronectin-binding proteins in *Staphylococcus aureus* are mediated by QS (Yarwood et al., 2004; Carnes et al., 2010). These QS-regulated virulence factors enhance the ability of pathogens to evade host immune defences, establish infections, acquire essential nutrients, and successfully colonize host tissues. Recently, quorum quenching agents, also referred to as anti-quorum sensing agents, have emerged as promising alternatives to conventional antibiotics due to their capacity to suppress bacterial virulence and enhance pathogen clearance in a variety of animal models. By targeting bacterial communication rather than bacterial survival, these agents can effectively prevent infection while minimizing the risk of resistance development. Although the clinical application of quorum quenching strategies remains in its early stages, growing evidence highlights their considerable therapeutic potential. This review aims to summarize the significance of quorum sensing signalling molecules and to discuss emerging strategies for preventing bacterial infections through interference with bacterial communication systems without promoting antimicrobial resistance.

Quorum Sensing Signals

Different classes of autoinducers are utilized by Gram-negative and Gram-positive bacteria to facilitate quorum sensing-mediated communication (LaSarre and Federle, 2013). In Gram-positive bacteria, the signalling molecules are predominantly autoinducing peptides (AIPs), whereas Gram-negative bacteria primarily employ N-acyl homoserine lactones (AHLs). Another important category of signalling molecules is Autoinducer-2 (AI-2), which is produced by both Gram-positive and Gram-negative bacteria and serves as a universal mediator of interspecies communication. These signalling compounds are released into the extracellular environment through diffusion and other transport mechanisms, enabling microorganisms to coordinate collective behaviours within both homogeneous and mixed microbial communities (Haque et al., 2019).

Molecular Mechanisms of Quorum Sensing

Quorum Sensing in Gram-Positive Bacteria

AIPs in gram-positive bacteria are recognized by membrane-bound two-component signal transduction systems, allowing bacterial populations to respond dynamically to changes in cell density within the extracellular environment (Sturme et al., 2002). AIPs originate from inactive precursor peptides known as pro-AIPs, which are synthesized intracellularly and subsequently processed into their active forms. Since these peptides cannot directly traverse the bacterial cell membrane, specialized transporter proteins are required for both their processing and export. Mature AIPs generally consist of 5–17 amino acids and may occur in either linear forms, as observed in *S. pneumoniae*, or cyclic structures, as found in *S. aureus*. Once released into the extracellular milieu, these peptides are detected by membrane-associated sensor kinases and initiates autophosphorylation of a conserved histidine residue within the receptor protein. The phosphate group is subsequently transferred to a cytoplasmic response regulator, which then activates the transcription of QS-responsive genes. Thus, the pro-AIP, transporter protein, histidine kinase receptor, and response regulator collectively function as an integrated signalling module that coordinates population-wide responses (Rutherford and Bassler, 2012).

The Accessory Gene Regulator (Agr) system of *S. aureus* serves as a classical example of quorum sensing in Gram-positive bacteria (Lyon et al., 2002; Murray and Williams, 2018). This regulatory network utilizes AIPs

as signalling molecules and controls the expression of numerous virulence determinants in response to increasing cell density. The principal components of the Agr system include AgrD, AgrB, AgrC, AgrA, the P2 and P3 promoters, and RNAIII. AgrD functions as the precursor peptide, which is processed and exported by the membrane-associated protein AgrB. As bacterial density increases, extracellular AIP concentrations rise and interact with the membrane-bound histidine kinase AgrC, triggering its autophosphorylation. The phosphate group is then transferred to the response regulator AgrA, which activates transcription from both the P2 and P3 promoters (Cheraghi et al., 2017). While the P2 promoter drives expression of the Agr operon itself, the P3 promoter facilitates the synthesis of RNAIII, a regulatory RNA molecule that orchestrates the expression of numerous virulence-associated genes (Paulander et al., 2018). Additional examples of QS systems in Gram-positive bacteria include the ComX/ComP/ComA regulatory system of *Bacillus subtilis*, where ComX functions as the signalling peptide (Spacapan et al., 2020), and the Fsr system of *Enterococcus faecalis*, which employs Gelatinase Biosynthesis Activating Pheromone (GBAP) as its signalling molecule (Ali et al., 2022).

Quorum Sensing in Gram-Negative Bacteria

In Gram-negative bacteria, quorum sensing is predominantly mediated through the LuxI/LuxR regulatory system, first identified in the bioluminescent marine bacterium *Vibrio fischeri* (Yang et al., 2009). This signalling mechanism regulates gene expression according to bacterial population density and relies on the synthesis and detection of AHL molecules. LuxI functions as an enzyme responsible for the synthesis of AHLs through reactions involving S-adenosylmethionine (SAM) and acyl carrier protein (ACP). As the bacterial population expands, AHL molecules accumulate in the extracellular environment and subsequently diffuse back into the cells, where they bind to the transcriptional regulator LuxR (Ivanova et al., 2018). In the absence of AHL, LuxR remains highly unstable and is rapidly degraded. However, the formation of the AHL–LuxR complex stabilizes the protein, enabling proper folding, DNA binding, and activation of target gene transcription. An important feature of this system is the positive feedback mechanism known as autoinduction. The AHL–LuxR complex activates transcription of the *luxI* gene, resulting in increased production of AHL molecules. This amplification loop facilitates rapid and synchronized activation of quorum sensing-controlled genes once a critical population density is achieved.

LuxI homologues possess distinct substrate-binding pockets that vary among bacterial species, enabling the synthesis of structurally diverse AHL molecules with acyl chain lengths ranging from C4 to C18. Additional modifications, including carbonyl and hydroxyl groups at the C3 position, further contribute to signal diversity. Correspondingly, LuxR homologues contain highly specific ligand-binding domains that recognize particular AHL structures. This remarkable chemical specificity ensures precise intraspecies communication and minimizes unintended cross-talk between different bacterial species (Papenfort and Bassler, 2016). In *Vibrio fischeri*, activation of the LuxI/LuxR circuit at high cell density stimulates transcription of the *luxICDABEG* operon, ultimately resulting in bioluminescence (Bellissimo et al., 2024). Other well-characterized quorum sensing systems in Gram-negative bacteria include LasI/LasR and RhlI/RhlR in *Pseudomonas aeruginosa* (Kostylev et al., 2019), SmaI/SmaR in *Serratia marcescens* (Van Houdt et al., 2007), CviI/CviR in *Chromobacterium violaceum* (Dimitrova et al., 2023), and VjbR, an orphan LuxR-family regulator, in *Brucella melitensis* (Uzureau et al., 2010).

Autoinducer-2 (AI-2)

Both Gram-positive and Gram-negative bacteria possess the capacity to detect and respond to interspecies communication signals through Autoinducer-2. Unlike AIPs and AHLs, which primarily mediate species-specific communication, AI-2 serves as a universal signalling molecule that facilitates interactions among diverse microbial populations. In the majority of bacterial species, AI-2 biosynthesis is catalysed by the enzyme LuxS (Thompson et al., 2015). Beyond its role in AI-2 production, LuxS constitutes a crucial component of the activated methyl cycle and exerts widespread regulatory effects on bacterial physiology. Studies have demonstrated that LuxS influences the expression of more than 400 genes associated with essential cellular functions, including surface adhesion, motility, biofilm formation, metabolic regulation, and toxin production (Hardie et al., 2003). Consequently, AI-2-mediated signalling is recognized as a central mechanism governing microbial community behaviour and interspecies coordination within complex ecological niches.

Role of Quorum Sensing in Bacterial Virulence

Quorum sensing enables bacterial populations to coordinate gene expression in a cell density-dependent manner, thereby facilitating collective behaviours that enhance survival, colonization, and pathogenicity. Through this highly organized communication network, bacteria ensure that the production of virulence-associated factors,

including toxins, extracellular enzymes, biofilms, motility systems, adhesion molecules, and immune evasion mechanisms, occurs only when the bacterial population has reached a sufficient density to exert a significant impact on the host (Riedel et al., 2001). Such coordinated regulation maximizes pathogenic efficiency while minimizing unnecessary metabolic expenditure. Among the numerous virulence determinants regulated by QS, toxins play a pivotal role in disrupting host barriers, suppressing immune responses, and facilitating access to essential nutrients. Similarly, extracellular enzymes such as proteases and lipases contribute to the degradation of host extracellular matrix components and cellular structures, thereby promoting tissue invasion and dissemination of infection. QS also regulates bacterial motility and chemotaxis, enabling microorganisms to respond dynamically to environmental stimuli, migrate toward favourable nutrient sources, and avoid adverse conditions. This adaptive behaviour enhances bacterial colonization and persistence within diverse ecological niches.

Surface adhesion molecules represent another critical group of QS-regulated factors. These structures mediate the initial attachment of bacteria to host tissues and abiotic surfaces, serving as a prerequisite for successful colonization and biofilm establishment. In addition, QS controls multiple immune evasion mechanisms that allow pathogens to circumvent host defence systems, suppress immune surveillance, and prolong bacterial survival within infected tissues. Biofilm formation is one of the most extensively studied QS-regulated processes. Biofilm development is a complex, multistage phenomenon involving initial attachment, cellular proliferation, microcolony formation, and the production of a self-generated extracellular polymeric matrix that ultimately gives rise to a highly organized three-dimensional architecture (Linnes et al., 2013). QS orchestrates not only the initiation and maturation of biofilms but also the eventual dispersal of bacterial cells from established biofilm communities. The biofilm matrix provides substantial protection against host immune responses and antimicrobial agents, making biofilm-associated infections particularly difficult to eradicate. Consequently, biofilm formation significantly contributes to the development and persistence of multidrug-resistant bacterial infections.

Quorum Sensing and Multidrug Resistance (MDR)

Although QS does not directly confer antibiotic resistance, it substantially enhances bacterial survival mechanisms that contribute to the emergence and persistence of MDR. Through the regulation of genes associated with biofilm formation, efflux pump activity, virulence factor production, stress adaptation, and horizontal gene transfer, QS creates a protective environment that reduces bacterial susceptibility to antimicrobial agents. The synchronized response induces a range of protective mechanisms that collectively enhance bacterial resilience against antimicrobial treatments, thereby making infections increasingly challenging to manage and eradicate.

Biofilm Formation

Quorum sensing does not directly interfere with biofilm formation; rather, it serves as a central regulatory system that promotes and coordinates biofilm development. Bacteria utilize QS signalling networks to synchronize the formation, maturation, and maintenance of biofilm communities in response to population density and environmental conditions. Biofilms function as highly effective physical and biochemical barriers that limit the penetration of antimicrobial agents and protect bacterial cells from host immune defenses. Bacteria embedded within biofilms can exhibit antimicrobial resistance levels up to 1,000-fold greater than those of their planktonic counterparts. By regulating the synthesis of extracellular matrix components and biofilm-associated genes, QS significantly enhances bacterial persistence and treatment resistance. For example, *Pseudomonas aeruginosa* employs the Las and Rhl quorum sensing systems to regulate key biofilm matrix constituents and maintain biofilm integrity (Williams and Camara, 2009).

Efflux Pumps

Quorum sensing and efflux pump systems are closely interconnected and collectively contribute to the development of MDR in microorganisms. The relationship between these systems is bidirectional: QS regulates the expression of efflux pumps, while efflux pumps can actively transport QS signalling molecules, thereby influencing communication pathways. QS-mediated upregulation of efflux pump genes results in the active extrusion of antibiotics from bacterial cells, reducing intracellular drug concentrations and diminishing antimicrobial efficacy. In *Pseudomonas aeruginosa*, the MexAB-OprM efflux pump system is regulated by QS and contributes significantly to resistance against fluoroquinolones, tetracyclines, and chloramphenicol (Maseda et al., 2004). The reciprocal interaction between QS and efflux pumps establishes a self-reinforcing cycle that promotes bacterial survival under antibiotic stress. Consequently, simultaneous inhibition of both systems has emerged as a promising therapeutic strategy for combating multidrug-resistant biofilm-associated infections.

Stress Response

Quorum sensing is intricately linked to bacterial stress response pathways and enables microorganisms to coordinate survival strategies under adverse environmental conditions. Beyond regulating collective behaviours, QS enhances bacterial tolerance to a variety of stressors, including antibiotic exposure, oxidative stress, osmotic stress, nutrient deprivation, and other environmental challenges (Lee et al., 2013). An important aspect of this adaptive response is the formation of persister cells, which are metabolically dormant bacterial subpopulations capable of surviving antimicrobial treatment. Although persister cells are not genetically resistant, their transient physiological state enables them to tolerate otherwise lethal concentrations of antibiotics. In *E. coli*, QS-regulated signalling pathways have been shown to promote persister cell formation under stressful conditions, thereby contributing to chronic and recurrent infections (Poole, 2012).

Horizontal Gene Transfer

Quorum sensing functions as an intelligent regulatory system for horizontal gene transfer, allowing bacterial populations to assess local cell density, environmental conditions, and the relative abundance of susceptible and resistant cells before initiating genetic exchange. This coordination ensures that energy-intensive transfer processes occur only when conditions are favourable and the probability of successful gene dissemination is high. By regulating conjugation, competence development, and DNA uptake mechanisms, QS enhances the transfer of antibiotic resistance genes and other adaptive traits between bacterial cells. Such regulation is particularly important within biofilms, where close cellular proximity facilitates efficient genetic exchange. QS promotes plasmid transfer and uptake of resistance determinants, including competence-mediated gene acquisition in *Streptococcus pneumoniae*, thereby accelerating the dissemination of antimicrobial resistance within bacterial populations (Van Gestel et al., 2021).

Regulation of Resistance Genes

Quorum sensing directly and indirectly regulates the expression, maintenance, and dissemination of antimicrobial resistance genes (ARGs) within bacterial communities. Certain resistance determinants are activated under high-cell-density conditions through QS-controlled regulatory pathways, ensuring that resistance mechanisms are expressed only when they provide a selective advantage. This regulation involves multiple interconnected mechanisms, including direct transcriptional activation of efflux pumps and β-lactamase genes, biofilm-mediated protection, modulation of reactive oxygen species (ROS), and enhancement of horizontal gene transfer. Through these coordinated pathways, QS optimizes energy utilization while maximizing bacterial survival under antimicrobial pressure. Consequently, targeting QS with specific inhibitors offers a promising approach to suppress both the expression and transmission of ARGs, thereby contributing to efforts aimed at mitigating antimicrobial resistance (Wang et al., 2023).

Quorum Quenching (QQ) and Quorum Sensing Inhibitors (QSIs)

Quorum quenching (QQ) represents an innovative anti-virulence strategy designed to attenuate bacterial pathogenicity and reduce resistance development by disrupting QS pathways rather than directly eliminating bacterial cells. Unlike conventional antibiotics, which impose strong selective pressure on microbial populations, QQ interferes with bacterial communication systems and thereby suppress the coordinated expression of virulence-associated traits. Quorum quenching agents inhibit cell-to-cell communication and interfere with QS-regulated activities such as toxin production, extracellular enzyme secretion, swimming, swarming, motility, biofilm formation, and host colonization. As a result, bacteria lose their ability to effectively establish infection and cause disease, while the likelihood of resistance-conferring mutations is reduced.

Several important characteristics should be considered when designing or selecting effective quorum sensing inhibitors (QSIs). Ideally, QSIs should be small, chemically stable molecules capable of efficiently suppressing QS-regulated gene expression. They should possess resistance to metabolic degradation and maintain biological activity for longer durations than native QS signals to ensure sustained inhibition. Furthermore, effective QSIs should selectively target pathogenic bacterial communication systems while preserving beneficial microbial populations that contribute to host health and microbial homeostasis. Quorum quenching agents encompass a diverse range of molecules, including QSIs, QQ peptides, QQ enzymes, cyclodextrins, and other bioactive compounds. Interestingly, certain conventional antibiotics, such as azithromycin, can also exhibit QQ activity when administered at sub-inhibitory concentrations (Hetta et al., 2024). QQ strategies include enzymatic degradation of signalling molecules, inhibition of signal synthesis and reception through small-molecule inhibitors, quorum signal mimicry and competitive antagonism, utilization of plant-derived anti-QS compounds, antibody-mediated signal neutralization, and active sequestration of autoinducer molecules by beneficial microorganisms.

Enzymatic Quorum Quenching

Enzymatic QQ involves the degradation or modification of signalling molecules responsible for bacterial communication. Key enzymes include AHL-lactonases and AHL-acylases, which hydrolyze the lactone ring and remove acyl side chains from AHL molecules, respectively (Uroz et al., 2008). Additionally, oxidoreductases can chemically modify autoinducers, converting them into inactive forms incapable of triggering QS responses.

Small-Molecule and Competitive Inhibitors

Small-molecule QSIs include halogenated furanones and a variety of naturally occurring compounds such as cinnamaldehyde, baicalin, ajoene, quercetin, and curcumin. These compounds interfere with diverse stages of QS pathways, including autoinducer synthesis, signal reception, receptor binding, and downstream signal transduction processes (Nazzaro et al., 2013). Certain natural and synthetic molecules function as structural mimics of native autoinducers and competitively inhibit receptor activation. In Gram-negative bacteria, halogenated furanones compete with AHL molecules for binding to LuxR-family receptors, thereby disrupting signal transmission (Wu et al., 2004). Similarly, in Gram-positive bacteria, synthetic peptide analogues can interfere with AIP receptor recognition and signalling *processes (Gong et al., 2024).*

Plant-Derived Quorum Sensing Inhibitors

Plant-derived phytochemicals have attracted considerable attention as natural QSIs due to their ability to suppress bacterial communication, virulence, and biofilm formation. Important examples include flavonoids such as quercetin and naringenin, terpenoids including thymol and carvacrol, and phenolic compounds such as eugenol and curcumin. These bioactive molecules target multiple components of QS pathways and represent promising candidates for the development of novel anti-virulence therapeutics.

Beneficial Bacteria as Quorum Quenching Agents

Certain beneficial bacterial strains can also contribute to quorum quenching by interfering with quorum sensing communication networks. Some strains of *E. coli*, including *E. coli* O157 and *E. coli* K12, are capable of sequestering signalling molecules and disrupting bacterial communication pathways through competitive interactions (Higuera-Ciapara et al., 2024). Such microbial interference may reduce pathogenic behaviour and contribute to maintaining ecological balance within microbial communities.

Quorum Quenching in Aquaculture

The rapid expansion and intensification of aquaculture practices have significantly increased the incidence and severity of bacterial diseases, particularly vibriosis, resulting in substantial economic losses worldwide. Growing concerns regarding antimicrobial resistance, coupled with increasingly stringent regulations governing antibiotic use in aquaculture, have stimulated the search for sustainable alternative disease-control strategies (Defoirdt et al., 2007). Among these, QQ has emerged as a promising approach for mitigating bacterial infections by disrupting QS-mediated pathogenicity rather than directly targeting bacterial viability. Several QQ-based strategies, including QSIs, naturally occurring algal metabolites, and signal-degrading microorganisms, have demonstrated considerable efficacy against pathogenic *Vibrio* species. Brominated furanones and their less toxic synthetic derivatives have been shown to interfere with AI-2-mediated QS pathways, thereby reducing bacterial virulence and enhancing the survival of aquatic organisms. Likewise, probiotic bacteria possessing AHL-degrading capabilities have successfully improved disease resistance and survival rates in fish and crustacean species by disrupting bacterial communication networks (Defoirdt et al., 2006).

Despite these encouraging outcomes, the ecological implications of QQ applications in aquaculture warrant careful consideration. QS not only regulates pathogenic behaviours but also influences beneficial microbial processes, including biofilm-mediated interactions that contribute to invertebrate larval settlement and ecosystem stability. Therefore, a comprehensive assessment of the long-term ecological impacts of QQ interventions is essential before their widespread implementation in aquaculture systems.

Quorum Quenching in Agriculture

Quorum quenching has emerged as an innovative and environmentally sustainable strategy for plant disease management by targeting QS-regulated virulence mechanisms rather than inhibiting bacterial growth directly. This anti-virulence approach minimizes selective pressure for resistance development while effectively reducing the pathogenic potential of phytopathogenic bacteria. Current QQ-based agricultural strategies involve the use of QQ-degrading enzymes and QSIs delivered through biocontrol microorganisms or transgenic plants engineered to express bacterial lactonases (D' Angelo Picard et al., 2011). Experimental studies have demonstrated significant reductions in disease severity

caused by QS-dependent phytopathogens such as *Pectobacterium*, while exerting minimal disruption on beneficial plant-associated microbial communities. Among naturally occurring quorum quenching microorganisms, soil-dwelling bacteria belonging to the genera *Bacillus* and *Rhodococcus* have attracted considerable attention because of their potent AHL-degrading activity and demonstrated biocontrol potential. These microorganisms offer promising opportunities for the development of sustainable agricultural practices aimed at reducing dependence on conventional chemical pesticides and antibiotics.

Quorum Quenching in Anti-Biofouling Applications

Bacterial biofilm formation is a major driving force behind biofouling in marine environments, industrial infrastructure, and water-treatment systems. Biofouling results in significant operational challenges, including increased maintenance costs, reduced system efficiency, accelerated material degradation, and environmental concerns. Conventional antifouling agents often rely on toxic chemicals that can adversely affect non-target organisms and aquatic ecosystems. Consequently, QQ has gained increasing attention as an environmentally friendly alternative that specifically targets QS-regulated biofilm development. By disrupting bacterial communication pathways, QQ strategies prevent the coordinated expression of genes required for biofilm formation and maturation.

Various QSIs derived from marine organisms and terrestrial plants, together with QS-degrading microorganisms and immobilized QQ enzymes, have demonstrated substantial efficacy in reducing biofilm formation under experimental conditions (Jiang et al., 2013). These findings highlight the potential of QQ-based technologies as sustainable and eco-friendly solutions for controlling biofouling across diverse industrial and environmental applications.

Quorum Quenching in Food Processing and Food Safety

Food safety and processing are profoundly influenced by the interactions among food-borne pathogens, spoilage microorganisms, and beneficial microbial communities. Many food-associated bacterial pathogens possess the ability to form persistent biofilms, a process that is frequently regulated by quorum sensing systems. These biofilms not only enhance bacterial survival under adverse conditions but also increase resistance to sanitation procedures and contribute to food contamination. QS-mediated biofilm formation plays a crucial role in the pathogenicity and persistence of several major food-borne pathogens, including *Salmonella*, *Listeria monocytogenes*, and *Vibrio parahaemolyticus*. The interplay between quorum sensing and biofilm development facilitates enhanced virulence, environmental adaptation, and persistence throughout food production and processing chains.

Numerous studies have demonstrated that QSIs, particularly brominated furanones, effectively disrupt AI-2 signalling pathways, suppress *luxS* gene expression, inhibit biofilm formation, and reduce extracellular enzyme production in these bacterial species (Phuvasate et al., 2012). Such findings underscore the considerable potential of quorum quenching strategies for minimizing biofilm-associated contamination, reducing pathogen persistence, and improving overall food safety standards within the food industry.

QSIs-Nanoparticles and QSIs Combined with Traditional Antibiotics

Recent advances in nanotechnology and antimicrobial therapeutics have opened new avenues for enhancing the efficacy of quorum sensing inhibitors. One promising approach involves the incorporation of QSIs into nanoparticle-based delivery systems or their strategic combination with conventional antibiotics through specialized carrier platforms (Han et al., 2017). Nanomaterials possess several advantageous properties, including large surface area-to-volume ratios, nanoscale dimensions, tunable physicochemical characteristics, and excellent biocompatibility. The integration of QSIs with nanomaterials can significantly improve therapeutic performance by enhancing compound stability, targeted delivery, controlled release, and bioavailability while simultaneously reducing systemic toxicity. Furthermore, combining QSIs with traditional antimicrobial agents offers multiple synergistic benefits. By suppressing QS-regulated virulence mechanisms, QSIs weaken bacterial defence and increase susceptibility to antibiotics. Such combination therapies can enhance antibacterial activity, promote selective bacterial eradication, reduce off-target toxicity, and achieve effective pathogen control using lower antibiotic dosages. Consequently, QSI-antibiotic combination strategies represent a promising approach for addressing multidrug-resistant bacterial infections and improving clinical treatment outcomes.

Challenges and Limitations of Quorum Quenching

Despite its considerable therapeutic promise, quorum quenching faces several scientific, technological, and regulatory challenges that must be addressed before widespread clinical and commercial adoption can be achieved.

i. One of the primary concerns associated with QQ is the potential for unintended off-target effects. Since many QQ agents lack strict species specificity, they may inadvertently interfere with beneficial microbiota or non-target bacterial quorum sensing systems, thereby altering microbial community dynamics.

ii. Prolonged exposure to QQ agents may drive the evolution of adaptive bacterial responses. Microorganisms can potentially develop compensatory mechanisms through mutations in QS receptor genes, overexpression of signal synthases, or activation of alternative regulatory pathways that bypass quorum sensing inhibition.

iii. Numerous naturally occurring QSIs, including flavonoids and garlic-derived compounds, exhibit poor aqueous solubility, limited chemical stability under physiological conditions, and inadequate tissue penetration. Consequently, advanced drug-delivery systems such as nanocarriers, hydrogels, liposomal formulations, and other innovative delivery platforms are required to enhance their bioavailability and therapeutic efficacy.

iv. Regulatory challenges also remain a significant obstacle. At present, clear and standardized regulatory frameworks governing the use of QSIs in livestock production, aquaculture, and other food-producing animal systems are limited. Establishing comprehensive safety and efficacy guidelines will be crucial for facilitating the practical implementation of QQ-based interventions.

Despite these limitations, quorum quenching continues to emerge as one of the most promising anti-virulence strategies in contemporary antimicrobial research. Ongoing advances in molecular biology, biotechnology, nanomedicine, and microbial ecology are expected to overcome current barriers and accelerate the translation of QQ-based approaches into next-generation therapeutic solutions for managing persistent and antimicrobial-resistant infections in both human and veterinary medicine.

Conclusions

In summary, QS represents far more than a simple bacterial communication mechanism; it functions as a master regulator governing numerous critical processes, including virulence factor production, biofilm development, antimicrobial resistance, stress adaptation, and horizontal gene transfer. Through the precise coordination of gene expression in response to population density, quorum sensing enables bacterial communities to optimize survival, colonization, and pathogenicity within diverse host and environmental settings. Targeting QS through QQ and other anti-virulence strategies offers a transformative paradigm in antimicrobial therapy. Rather than exerting bactericidal pressure, these approaches selectively disarm pathogens by disrupting communication networks that regulate disease-causing behaviours. Such interventions reduce the selective pressures that drive resistance development, restore the effectiveness of existing antibiotics against multidrug-resistant biofilm-associated infections, and limit the dissemination of antimicrobial resistance genes by interfering with conjugation and horizontal gene transfer mechanisms.

Furthermore, QQ strategies can act synergistically with conventional antibiotics, enhancing therapeutic efficacy while reducing the dosage requirements of traditional antimicrobial agents. As the global burden of antimicrobial resistance continues to escalate, QS-targeted interventions represent a sustainable and potentially low-resistance alternative to conventional antimicrobial therapies. Future research focused on optimizing QSIs, improving targeted delivery systems, understanding ecological impacts, and establishing robust regulatory frameworks will be essential for translating these promising discoveries into practical clinical applications. Ultimately, the transition from strategies aimed at "killing" bacteria to those designed to "disarm" them may redefine the future of infectious disease management and provide a powerful weapon in the ongoing battle against antimicrobial resistance.

Conflicts of interest and financial disclosures

The authors state that there are no conflicts of interest to disclose.

References

Ali, I. A. A., Levesque, C. M., & Neelakantan, P. (2022). Fsr quorum sensing system modulates the temporal development of Enterococcus faecalis biofilm matrix. *Molecular Oral Microbiology, 37*(1), 22–30.

Antunes, L. C. M., Ferreira, R. B. R., Buckner, M. M. C., & Finlay, B. B. (2010). Quorum sensing in bacterial virulence. *Microbiology, 156*(8), 2271–2282.

Bellissimo, K. A., Septer, A. N., Whistler, C. A., Rodríguez, C., & Stabb, E. V. (2024). Deletion of luxI increases luminescence of Vibrio fischeri. mBio, 15(10), e0244624.

Bramhachari, P. V., Yugandhar, N. M., & Prathyusha, A. M. V. N. (2018). Quorum sensing regulated swarming motility and migratory behavior in bacteria. In P. V. Bramhachari (Ed.), Implication of quorum sensing system in biofilm formation and virulence (pp. 49–66).

Carnes, E. C., Lopez, D. M., & Donegan, N. P. (2010). Confinement induced quorum sensing of individual Staphylococcus aureus bacteria. *Nature Chemical Biology, 6*(1), 41–45.

Cheraghi, S., Pourgholi, L., & Shafaati, M. (2017). Analysis of virulence genes and accessory gene regulator (agr) types among methicillin-resistant Staphylococcus aureus strains in Iran. *Journal of Global Antimicrobial Resistance, 10*, 315–320.

D'Angelo Picard, C., Chapelle, E., & Ratet, P. (2011). Transgenic plants expressing the quorum quenching lactonase AttM do not significantly alter root-associated bacterial populations. *Research in Microbiology, 162*(1), 951–958.

Defoirdt, T., Boon, N., & Sorgeloos, P. (2007). Alternatives to antibiotics to control bacterial infections: Luminescent vibriosis in aquaculture as an example. *Trends in Biotechnology, 25*(10), 472–479.

Defoirdt, T., Crab, R., Wood, T. K., Sorgeloos, P., Verstraete, W., & Bossier, P. (2006). Quorum sensing-disrupting brominated furanones protect the gnotobiotic brine shrimp Artemia franciscana from pathogenic Vibrio harveyi, Vibrio campbellii, and Vibrio parahaemolyticus isolates. *Applied and Environmental Microbiology, 72*(9), 6419–6423.

Dimitrova, P. D., Damyanova, T., & Paunova Krasteva, T. (2023). Chromobacterium violaceum: A model for evaluating the anti-quorum sensing activities of plant substances. *Scientia Pharmaceutica, 91*(3), 33.

Gong, X., Brand, C. J., & Bertucci, M. A. (2024). Designing and synthesizing peptide-based quorum sensing modulators. *Methods in Enzymology, 698*, 263–299.

Han, M., Gu, J., Gao, G. F., & Liu, W. J. (2017). China in action: National strategies to combat emerging infectious diseases. *Science China Life Sciences, 60*(12), 1383–1385.

Haque, S., Yadav, D. K., & Bisht, S. C. (2019). Quorum sensing pathways in Gram-positive and Gram-negative bacteria: Potential of their interruption in abating drug resistance. *Journal of Chemotherapy, 31*(4), 161–187.

Hardie, K. R. (2003). Autoinducer-2 activity in Escherichia coli culture supernatants can be actively reduced despite maintenance of an active synthase, LuxS. *Microbiology, 149*(3), 715–728.

Hetta, H. F., Ramadan, Y. N., Rashed, Z. I., Alharbi, A. A., Alsharef, S., Alkindy, T. T., Alkhamali, A., Albalawi, A. S., Battah, B., & Donadu, M. G. (2024). Quorum sensing inhibitors: An alternative strategy to win the battle against multidrug-resistant (MDR) bacteria. *Molecules, 29*(15), 3466.

Higuera-Ciapara, I., Benitez-Vindiola, M., Figueroa-Yañez, L. J., & Martínez-Benavidez, E. (2024). Polyphenols and CRISPR as quorum quenching agents in antibiotic-resistant foodborne human pathogens (Salmonella Typhimurium, Campylobacter jejuni and Escherichia coli O157:H7). *Foods, 13*(4), 584.

Ivanova, A., Ivanova, K., & Tzanov, T. (2018). Inhibition of quorum sensing: A new paradigm in controlling bacterial virulence and biofilm formation. In V. C. Kalia (Ed.), Biotechnological applications of quorum sensing inhibitors (pp. 5–10).

Jiang, W., Xia, S., Liang, J., Zhang, Z., & Hermanowicz, S. W. (2013). Effect of quorum quenching on reactor performance, biofouling and biomass characteristics in membrane bioreactors. *Water Research, 47*(1), 187–196.

Kostylev, M., Kim, D. Y., & Smalley, N. E. (2019). Evolution of the Pseudomonas aeruginosa quorum sensing hierarchy. *Proceedings of the National Academy of Sciences, 116*(14), 7027–7032.

LaSarre, B., & Federle, M. J. (2013). Exploiting quorum sensing to confuse bacterial pathogens. *Microbiology and Molecular Biology Reviews, 77*(1), 73–111.

Lee, J., Wu, J., Deng, Y., Wang, J., Wang, C., Wang, J., Chang, C., Dong, Y., Williams, P., & Zhang, L. H. (2013). A cell-cell communication signal integrates quorum sensing and stress response. *Nature Chemical Biology, 9*(5), 339–343.

Lewis, K. (2007). Persister cells, dormancy and infectious disease. *Nature Reviews Microbiology, 5*(1), 48–56.

Li, J., Chen, J., Vidal, J. E., & McClane, B. A. (2011). The Agr-like quorum-sensing system regulates sporulation and production of enterotoxin and Beta2 toxin by Clostridium perfringens type A non-food-borne human gastrointestinal disease strain F5603. *Infection and Immunity, 79*(6), 2451–2459.

Linnes, J. C., Ma, H., & Bryers, J. D. (2013). Giant extracellular matrix binding protein expression in Staphylococcus epidermidis is regulated by biofilm formation and osmotic pressure. *Current Microbiology, 66*(6), 627–633.

Lyon, G. J., Wright, J. S., Muir, T. W., & Novick, R. P. (2002). Key determinants of receptor activation in the agr autoinducing peptides of Staphylococcus aureus. *Biochemistry, 41*(31), 10095–10104.

Maseda, H., Sawada, I., Saito, K., Uchiyama, H., Nakae, T., & Nomura, N. (2004). Enhancement of the mexAB-oprM efflux pump expression by a quorum sensing autoinducer and its cancellation by MexT. *Antimicrobial Agents and Chemotherapy, 48*(4), 1320–1328.

Murray, E. J., & Williams, P. (2018). Detection of agr-type autoinducing peptides produced by Staphylococcus aureus. In Methods in Molecular Biology (Vol. 1673, pp. 89–96).

Musk, D., Jr., & Hergenrother, P. (2006). Chemical countermeasures for the control of bacterial biofilms: Effective compounds and promising targets. *Current Medicinal Chemistry, 13*(18), 2163–2177.

Nazzaro, F., Fratianni, F., & Coppola, R. (2013). Quorum sensing and phytochemicals. *International Journal of Molecular Sciences, 14*(6), 12607–12619.

Papenfort, K., & Bassler, B. L. (2016). Quorum sensing signal-response systems in Gram-negative bacteria. *Nature Reviews Microbiology, 14*(9), 576–588.

Paulander, W., Varming, A. N., Bojer, M. S., Friberg, C., Bæk, K., & Ingmer, H. (2018). The agr quorum sensing system in Staphylococcus aureus cells mediates death of sub-population. *BMC Research Notes, 11*(1), 503.

Phuvasate, S., Chen, M. H., & Su, Y. C. (2012). Reductions of Vibrio parahaemolyticus in Pacific oysters by depuration at various temperatures. *Food Microbiology, 31*(1), 166–170.

Poole, K. (2012). Bacterial stress responses as determinants of antimicrobial resistance. *Journal of Antimicrobial Chemotherapy, 67*(9), 2069–2089.

Popham, D. L., & Stevens, A. M. (2006). Bacterial quorum sensing and bioluminescence. *Tested Studies for Laboratory Teaching, 27*, 201–215. Association for Biology Laboratory Education.

Preda, V. G., & Sandulescu, O. (2019). Communication is the key: Biofilms, quorum sensing, formation and prevention. *Discoveries, 7*, e100.

Riedel, K., Gotschlich, A., & Givskov, M. (2001). The cep quorum sensing system of Burkholderia cepacia H111 controls biofilm formation and swarming motility. *Microbiology, 147*(9), 2517–2528.

Rutherford, S. T., & Bassler, B. L. (2012). Bacterial quorum sensing: Its role in virulence and possibilities for its control. *Cold Spring Harbor Perspectives in Medicine, 2*(11), a012427.

Spacapan, M., Danevčič, T., Štefanič, P., Porter, M., Stanley-Wall, N. R., & Mandic-Mulec, I. (2020). The ComX quorum sensing peptide of Bacillus subtilis affects biofilm formation negatively and sporulation positively. *Microorganisms, 8*(8), 1131.

Sturme, M. H., Kleerebezem, M., Nakayama, J., Akkermans, A. D., Vaughan, E. E., & De Vos, W. M. (2002). Cell-to-cell communication by autoinducing peptides in Gram-positive bacteria. *Antonie van Leeuwenhoek, 81*(1–4), 233–243.

Thompson, J., Oliveira, R., Djukovic, A., Ubeda, C., & Xavier, K. (2015). Manipulation of the quorum sensing signal AI-2 affects the antibiotic-treated gut microbiota. *Cell Reports, 10*(11), 1861–1871.

Uroz, S., Oger, P. M., Chapelle, E., Adeline, M. T., Faure, D., & Dessaux, Y. (2008). A Rhodococcus qsdA-encoded enzyme defines a novel class of large-spectrum quorum-quenching lactonases. *Applied and Environmental Microbiology, 74*(5), 1357–1366.

Uzureau, S., Lemaire, J., & Delaive, E. (2010). Global analysis of quorum sensing targets in the intracellular pathogen Brucella melitensis 16M. *Journal of Proteome Research, 9*(6), 3200–3217.

Van Gestel, J., Bareia, T., Tenennbaum, B., Dal Co, A., Guler, P., Aframian, N., Puyesky, S., Grinberg, I., D'Souza, G. G., Erez, Z., Ackermann, M., & Eldar, A. (2021). Short-range quorum sensing controls horizontal gene transfer at micron scale in bacterial communities. *Nature Communications, 12*(1), 2324.

Van Houdt, R., Givskov, M., & Michiels, C. W. (2007). Quorum sensing in Serratia. *FEMS Microbiology Reviews, 31*(4), 407–424.

Wang, M., Lian, Y., Wang, Y., & Zhu, L. (2023). The role and mechanism of quorum sensing on environmental antimicrobial resistance. *Environmental Pollution, 322*, 121238.

Williams, P., & Camara, M. (2009). Quorum sensing and environmental adaptation in Pseudomonas aeruginosa: A tale of regulatory networks and multifunctional signal molecules. *Current Opinion in Microbiology, 12*(2), 182–191.

Wu, H., Song, Z., Hentzer, M., Andersen, J. B., Molin, S., Givskov, M., & Høiby, N. (2004). Synthetic furanones inhibit quorum-sensing and enhance bacterial clearance in Pseudomonas aeruginosa lung infection in mice. *The Journal of Antimicrobial Chemotherapy, 53*(6), 1054–1061.

Yang, M., Sun, K., Zhou, L., Yang, R., Zhong, Z., & Zhu, J. (2009). Functional analysis of three AHL autoinducer synthase genes in Mesorhizobium loti reveals the important role of quorum sensing in symbiotic nodulation. *Canadian Journal of Microbiology, 55*(2), 210–214.

Yarwood, J. M., Bartels, D. J., Volper, E. M., & Greenberg, E. P. (2004). Quorum sensing in Staphylococcus aureus biofilms. *Journal of Bacteriology, 186*(6), 1838–1850.